

Immunisation Schedule Queensland Children (under 10 years)

April 2019

Before vaccinating:

- **Review** the **Australian Immunisation Register (AIR)** to check the child's previous immunisation history
- Check the **online Australian Immunisation Handbook** for information about reconstitution of vaccines, site to administer, catch-up vaccination, vaccination for special risk groups, etc at <https://immunisationhandbook.health.gov.au/>
- **Apply sequential dose numbers** for each vaccine administered and **report all records** immediately to **AIR**

Age	Disease	Vaccine brand	Important notes
Birth	Hepatitis B	H-B-VaxII paediatric OR Engerix B paediatric	<ul style="list-style-type: none"> • Give within 24 hours of birth • Can be given up to 7 days after birth
	Tuberculosis	BCG	<ul style="list-style-type: none"> • Aboriginal & Torres Strait Islander children *Please see BCG note over the page
2 months (can be given from 6 weeks) and 4 months	DTPa-hepB-IPV-Hib	Infanrix Hexa - <i>reconstitute</i>	ALL CHILDREN <ul style="list-style-type: none"> • 1st dose must be given <15 weeks of age • 2nd dose must be given <25 weeks of age Check the Rotarix wheel for timing of vaccination
	Pneumococcal	Prevenar 13	
	Rotavirus	Rotarix	
6 months	DTPa-hep B-IPV-Hib	Infanrix Hexa - <i>reconstitute</i>	ALL CHILDREN <ul style="list-style-type: none"> • Aboriginal & Torres Strait Islander children • Children with medical risk factors • Premature infants born <28 weeks gestation
	Pneumococcal	Prevenar 13	
From 6 months to <5 years	Influenza	As supplied	ALL CHILDREN should receive influenza vaccine annually <ul style="list-style-type: none"> • 2 doses required for children from 6 months to <9 years of age with a minimum of 1 month between doses in the first year they receive influenza vaccine Please refer to the Influenza Program Advice



Age	Disease	Vaccine brand	Important notes
12 months	Measles-mumps-rubella	Priorix - <i>reconstitute</i> OR MMRII - <i>reconstitute</i>	ALL CHILDREN
	Meningococcal ACWY	Nimenrix	
	Pneumococcal	Prevenar 13	
	Hepatitis A	Vaqa paediatric	Aboriginal & Torres Strait Islander children
	Hepatitis B	H-B-VaxII paediatric OR Engerix B paediatric	Premature baby <32 weeks gestation or <2000g birthweight only
18 months	Measles-mumps-rubella- varicella	Priorix Tetra - <i>reconstitute</i> OR Proquad - <i>reconstitute</i>	ALL CHILDREN
	<i>Haemophilus influenzae</i> type b	Act-HIB - <i>reconstitute</i>	ALL CHILDREN
	DTPa	Infanrix OR Tripacel	ALL CHILDREN
	Hepatitis A	Vaqa paediatric	Aboriginal & Torres Strait Islander children
4 years	DTPa-IPV	Infanrix IPV OR Quadracel	ALL CHILDREN
	Pneumococcal	Pneumovax 23	1 dose for children with medical risk factors
5 years and older	Influenza	As supplied	ALL Aboriginal & Torres Strait Islander children from 6 months of age

*BCG is provided by TB Control Units and is recommended for:

- Aboriginal and Torres Strait Islander newborns and other children <5 years of age living in Indigenous communities
- Children <5 years travelling to a country of high TB incidence for a prolonged period
- Newborns where there is a family history of leprosy

Further information can be obtained from www.health.qld.gov.au/__data/assets/pdf_file/0027/637614/bcg-English.pdf

